arXiv:0912.3471v1 [math.MG] 17 Dec 2009

Isometries of Products of Path-Connected Locally Uniquely Geodesic Metric Spaces with the Sup Metric are Reducible.

William Malone

December 17, 2009

Abstract

Let M_i and N_i be path-connected locally uniquely geodesic metric spaces that are not points and $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ be an isometry where $\prod_{i=1}^n N_i$ and $\prod_{i=1}^m M_i$ are given the sup metric. Then $m = n$ and after reindexing M_i is isometric to N_i for all i. Moreover f is a composition of an isometry that reindexes the factor spaces and an isometry that is a product of isometries $f_i : M_i \to N_i$.

1 Introduction

For the duration a product $\prod_{i=1}^m M_i$ of metric spaces is always considered to be endowed with the sup metric. In these metric spaces there are two obvious types of isometries $f : \prod_{i=1}^m M_i \to \prod_{i=1}^m M_i$. The first is a product $f = f_1 \times \cdots \times f_m$ where $f_j : M_j \to M_j$ is an isometry. The second is a reindexing of the form

$$g(\alpha_1, \ldots, \alpha_m) = (\alpha_{\pi^{-1}(1)}, \ldots, \alpha_{\pi^{-1}(m)})$$

for some permutation π of $\{1, \ldots, m\}$. This leads to the following definition.

Definition 1.1. *An isometry $f : \prod_{i=1}^m M_i \to \prod_{i=1}^m N_i$ is reducible if there is a permutation π of $\{1, 2, \ldots, m\}$ and isometries $f_i : M_i \to N_{\pi(i)}$ such that*

$$f(\alpha_1, \ldots, \alpha_m) = (f_{\pi^{-1}(1)}(\alpha_{\pi^{-1}(1)}), \ldots, f_{\pi^{-1}(m)}(\alpha_{\pi^{-1}(m)})).$$

The main result states that in certain cases all isometries are reducible.

Main Theorem. *Suppose M_i and N_i are path-connected locally uniquely geodesic metric spaces that are not points with $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ an isometry. Then $m = n$ and f is reducible.*

The heart of the proof lies in the fact that particular isometric embeddings of a graph in the product space $\prod_{i=1}^m M_i$ are invariant under any isometry. This will quickly lead to the number of factors in the product being an isometry invariant as well as forcing f to take the aspect of a product map. A proof for the case where $m = 2$ occurs in the appendix of [2], and a discussion of the background on geodesic metric spaces can be found in [1].

1.1 Acknowledgements

I would like to thank Mladen Bestvina, Yael Algom-Kfir, and Erika Meucci for their many helpful comments and discussions. I would also like to thank Benson Farb for mentioning this problem during a talk at the University of Utah.

2 Proof of Main Theorem

In this section we define a useful family of subsets of $\prod_{i=1}^m M_i$. Utilizing the Main Lemma, a proof of the Main Theorem is then provided.

Definition 2.1. *A k-slice Ω is a subset of $\prod_{i=1}^m M_i$ of the form*

$$\alpha_1 \times \cdots \times \alpha_{k-1} \times \Omega_k \times \alpha_{k+1} \times \cdots \times \alpha_m$$

where $\alpha_i \in M_i$ and $\Omega_k \subset M_k$ with $|\Omega_k| \geq 2$. The α_i are called the fixed coordinates of Ω.

Main Lemma. *Suppose that M_i and N_i are path-connected locally uniquely geodesic metric spaces and $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ an isometry. If $\Omega \subset \prod_{i=1}^m M_i$ is a k-slice then $f(\Omega)$ is a j-slice. Moreover Ω and Ω' are both k-slices if and only if $f(\Omega)$ and $f(\Omega')$ are both j-slices.*

In order to prove the Main Lemma and hence the Main Theorem the fact that $m = n$ for any isometry $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ is essential. However the proof of this fact is more technical and will be postponed.

Proof. (of Main Theorem) Assuming $m = n$, let $\Phi \subset \prod_{i=1}^m M_i$ be a k-slice for $k \in \{1, \ldots, m\}$. By the Main Lemma the image $f(\Phi)$ is a j-slice for some j. The second part of the Main Lemma gives more information namely that the image of any k-slice $\Omega \subset \prod_{i=1}^m M_i$ will be a j-slice.

To complete the proof it suffices to show that if two points,

$$a = (\alpha_1, \alpha_2, \ldots, \alpha_{k-1}, \gamma, \alpha_{k+1}, \ldots, \alpha_m)$$

and

$$b = (\beta_1, \beta_2, \ldots, \beta_{k-1}, \gamma, \beta_{k+1}, \ldots, \beta_m)$$

such that $\alpha_i \neq \beta_i$ for all i, have the same k^{th} coordinate then $f(b)$ and $f(a)$ have the same j^{th} coordinate. This will show the existence of isometries $f_i : M_k \to N_j$ and complete the proof of reducibility after post composing by the reindexing isometry

$\pi : \prod_{i=1}^m N_i \to \prod_{i=1}^m N_i$ where $\pi(N_j) = N_k$. Consider the points

$$\begin{aligned}
&a_0 = a = (\alpha_1, \alpha_2, \ldots, \alpha_{k-1}, \gamma, \alpha_{k+1}, \ldots, \alpha_m)\\
&a_1 = (\beta_1, \alpha_2, \ldots, \alpha_{k-1}, \gamma, \alpha_{k+1}, \ldots, \alpha_m)\\
&a_2 = (\beta_1, \beta_2, \alpha_3, \ldots, \alpha_{k-1}, \gamma, \alpha_{k+1}, \ldots, \alpha_m)\\
&\vdots\\
&a_{k-1} = (\beta_1, \beta_2, \ldots, \beta_{k-1}, \gamma, \alpha_{k+1}, \ldots, \alpha_m)\\
&a_k = (\beta_1, \beta_2, \ldots, \beta_{k-1}, \gamma, \beta_{k+1}, \alpha_{k+2}, \ldots, \alpha_m)\\
&\vdots\\
&a_{m-2} = (\beta_1, \beta_2, \ldots, \beta_{k-1}, \gamma, \beta_{k+1}, \ldots, \beta_{m-1}, \alpha_m)\\
&a_{m-1} = b = (\beta_1, \beta_2, \ldots, \beta_{k-1}, \gamma, \beta_{k+1}, \ldots, \beta_m).
\end{aligned}$$

Notice that $\{a_i, a_{i+1}\}$ is an $i+1$-slice for $i \in \{0, 1, \ldots, k-2\}$ and an $i+2$ slice for $i \in \{k-1, \ldots, m-2\}$. Since no pair $\{a_i, a_{i+1}\}$ is a k-slice, the pair of points $\{f(a_i), f(a_{i+1})\}$ is never a j-slice by the Main Lemma. Thus the j^{th} coordinate of $f(a_i)$ and $f(a_{i+1})$ are identical for all i. $\square$

3 The Graph $\mathcal{Q}_r^m$

In this section we define a metric graph, give examples of particularly nice isometric embeddings of this graph in $\prod_{i=1}^m M_i$, show that $m = n$ for any isometry $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$, and prove the Main Lemma.

Observe that if $\alpha_i, \beta_i \in M_i$ are connected by a unique geodesic of length r for $i = 1, \ldots, m$ then $a = (\alpha_1, \ldots, \alpha_m)$ and $b = (\beta_1, \ldots, \beta_m)$ are connected by a unique geodesic in $\prod M_i$. This fact that in certain directions within the product pairs of points are joined by unique geodesics is the main motivation for the following graph.

Definition 3.1. *A quadrilateral graph of dimension m with length r denoted $\mathcal{Q}_r^m \subset \prod_{i=1}^m \mathbb{R}$ is the metric graph whose vertices are of the form $\pm e_i^r = (0, 0, \ldots, 0, \pm 2r, 0, \ldots, 0)$ (non-zero in i^{th} coordinate) or $(\pm r, \pm r, \ldots, \pm r)$. Connect two vertices by an edge of length r if their distance is r in $\prod_{i=1}^m \mathbb{R}$.*

The first two examples of this family of graphs (pictured in Figure 1) are a quadrilateral with subdivided sides and the one skeleton of Kepler's Rhombic dodecahedron [3].

Example 3.2. *Let M_i be a uniquely geodesic metric space. An example of an isometric embedding $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ is given by the following construction. Pick a geodesic segment $\lambda_i \subset M_i$ of length $4r$. Let $\lambda_i(0) = \alpha_i$, $\lambda_i(r) = \theta_i$ $\lambda_i(2r) = \beta_i$, $\lambda_i(3r) = \varphi_i$ and $\lambda_i(4r) = \omega_i$. Then let*

$$\begin{aligned}
&\iota(e_k^r) = \{(\beta_1, \beta_2, \ldots, \beta_{k-1}, \omega_k, \beta_{k+1}, \ldots, \beta_m \| k \in \{1, 2, \ldots, m\}\}\\
&\iota(-e_k^r) = \{(\beta_1, \beta_2, \ldots, \beta_{k-1}, \alpha_k, \beta_{k+1}, \ldots, \beta_m \| k \in \{1, 2, \ldots, m\}\}\\
\text{and as a set}\quad &\iota(\pm r, \pm r, \ldots, \pm r) = \{(\pi_1, \ldots, \pi_m) | \pi_i \in \{\theta_i, \varphi_i\}\}
\end{aligned}$$

which defines the isometry on the set of vertices. One can check that appropriate vertices are connected by unique geodesics of length r.



Figure 1: $\mathcal{Q}_1^3 \subset \prod_{i=1}^3 \mathbb{R}$ (left) and $\mathcal{Q}_1^2 \subset \prod_{i=1}^2 \mathbb{R}$ (right)

This construction shows that isometric embeddings $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ are easy to construct when the M_i are uniquely geodesic metric spaces. If the spaces M_i are only path-connected locally uniquely geodesic metric spaces then in general an isometric embedding $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ will not exist for large r. However for every $a \in \prod M_i$ there exists a metric ball $B_\epsilon(a)$ such that $B_\epsilon(a)$ is a product of uniquely geodesic metric spaces. Thus for $4r < \epsilon$ an isometrically embedded $\mathcal{Q}_r^m$ can be constructed in the same way.

Definition 3.3. *Let $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ be an isometric embedding. If the image of every edge is a uniquely geodesic segment in $\prod M_i$ then call $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ admissible.*

If $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ is admissible and $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ is an isometry then $f \circ \iota : \mathcal{Q}_r^k \to \prod_{i=1}^n N_i$ is admissible. This is clear from definitions.

Definition 3.4. *Let $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ be admissible. The map ι is standard if for all j there exists an l such that $\{\iota(e_j^r), \iota(-e_j^r)\}$ is an $l-$slice.*

The proof of the Main Lemma requires a result establishing the fact that every admissible isometric embedding $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ is standard. The following discussion works toward establishing this fact. First some results concerning the behavior of points in the image of an admissible $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ are required.

Property 1. *For any vertex $x_j \in Q_r^m$ of the form $\{\pm e_1^r, \ldots, \pm e_m^r\}$ and any pair of points $\{e_i^r, -e_i^r\}$ with $x_j \notin \{e_i^r, -e_i^r\}$ there exists a geodesic segment $\Lambda \subset \mathcal{Q}_r^m$ with endpoints $\{e_i^r, -e_i^r\}$ such that $x_j \in \Lambda$.*

Constructing such a piecewise uniquely geodesic path in $\mathcal{Q}_r^m$ is straightforward. Note that there are precisely 2^{2m-4} paths connecting pairs of endpoints $\{e_i^r, -e_i^r\}$ so Λ is unique only when $m = 2$.

Property 2. *Assume that M_i is a uniquely geodesic metric space. Let p and q be points in $\prod_{i=1}^m M_i$ and Λ a geodesic with endpoints $\{p, q\}$. Let $d_{M_i}(p_i, q_i)$ be the distance between the i^{th} coordinates of p and q. Note that $d_{M_i}(p_i, q_i) = d(p, q)$ if and only if the i^{th} coordinate path of Λ (denoted λ_i) is a geodesic of length $d(p, q)$. Moreover for any point $z \in \Lambda$ the i^{th} coordinate of z is uniquely determined by $d(p, z)$ (since λ_i is the only geodesic connecting its endpoints in M_i).*

Property 2 follows directly from the definition of a geodesic and will allow us to characterize the possible admissible $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ when M_i is uniquely geodesic.

Lemma 3.5. *Assume that M_i is a uniquely geodesic metric space and let $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ be admissible. Let*

$$\begin{array}{c} \iota(e_j^r) = (\alpha_1, \ldots, \alpha_m) \\ \iota(-e_j^r) = (\beta_1, \ldots, \beta_m) \\ \iota(e_t^r) = (\gamma_1, \ldots, \gamma_m) \\ \iota(-e_t^r) = (\kappa_1, \ldots, \kappa_m) \end{array} .$$

If $d(\iota(e_t^r), \iota(-e_t^r)) = d_{M_l}(\gamma_l, \kappa_l) = 4r$ then $\alpha_l = \beta_l$.

Proof. By property 1 there exists geodesics $\Lambda, \Lambda' \subset \iota(\mathcal{Q}_r^k)$ such that the endpoints of Λ and Λ' are $\iota(\pm e_t^r)$ with $\iota(e_j^r) \in \Lambda$ and $\iota(-e_j^r) \subset \Lambda'$. By property 2 and the fact that $d(\iota(e_t^r), \iota(e_j^r)) = d(\iota(e_t^r), \iota(-e_j^r)) = 2r$ we see that $\alpha_l = \beta_l$. □

For an admissible $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ associate a number

$$q_j = |\{l \in \{1, \ldots, m\} | d(\iota(e_j^r), \iota(-e_j^r)) = d_{M_l}(\alpha_l, \beta_l)\}|$$

to each pair $\iota(\pm e_j^r)$. Clearly $q_j \geq 1$ for all j. If follows immediately from Lemma 3.5 that two pairs $\{\iota(e_j^r), \iota(-e_j^r)\}$ and $\{\iota(e_t^r), \iota(-e_t^r)\}$ cannot attain the distance $4r$ in the same coordinate. These two observations imply $q_j = 1$ for all j when $k = m$.

Lemma 3.6. *Let M_i and N_i be path-connected locally uniquely geodesic metric spaces and $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ be an isometry. Then $m = n$.*

Proof. (Of lemma 3.6) By invariance of an admissible $\iota(\mathcal{Q}_r^k)$ under isometry it follows that the largest k such that there exists an admissible $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ is an isometry invariant. Let this number be denoted by $L(\prod_{i=1}^m M_i)$. There exists a ball $B(a, 4r)$ around any point $a \in \prod_{i=1}^m M_i$ such that $B(a, 4r)$ is a product of uniquely geodesic metric spaces. An admissible $\mathcal{Q}_r^m$ can be constructed in $B(a, 4r)$ by Example 3.2 which shows $L(\prod_{i=1}^m M_i) \geq m$. Since the number $q_j \geq 1$ for all pairs $\{\iota(e_j^r), \iota(-e_j^r)\}$ in any admissible $\iota : \mathcal{Q}_r^k \to \prod_{i=1}^m M_i$ and $\sum_{j=1}^m q_j \leq m$ it is obvious that for $k > m$ no such ι can exist and thus $L(\prod_{i=1}^m M_i) \leq m$. □

Lemma 3.7. *Let M_i be a uniquely geodesic metric space and $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ admissible. Then $\iota : \mathcal{Q}_r^m \to \prod_{i=1}^m M_i$ is standard.*

Proof. (Of Lemma 3.7) Using the above fact that $q_j = 1$ for all j and Lemma 3.5 the collection of pairs

$$\left(\cup_{i=1}^{j-1}\{\iota(e_i^r), \iota(-e_i^r)\}\right) \cup \left(\cup_{i=j+1}^{n}\{\iota(e_i^r), \iota(-e_i^r)\}\right)$$

forces the distance between coordinates of $\{\iota(e_j^r), \iota(-e_j^r)\}$ to be either $4r$ or 0, thus making $\{\iota(e_j^r), \iota(-e_j^r)\}$ a l-slice for some l. □

We now restate and prove the Main Lemma.

Main Lemma. *Suppose that M_i and N_i are path-connected locally uniquely geodesic metric spaces and $f : \prod_{i=1}^m M_i \to \prod_{i=1}^n N_i$ an isometry. If $\Omega \subset \prod_{i=1}^m M_i$ is a k-slice then $f(\Omega)$ is a j-slice. Moreover Ω and Ω' are both k-slices if and only if $f(\Omega)$ and $f(\Omega')$ are both j-slices.*

Proof. By lemma 3.6 $m = n$. Three things must be shown.

Step 1 (Part A) Given three distinct points a, b, c such that $\{a, b\}$ is a j-slice, $\{b, c\}$ is a k-slice, and $\{a, c\}$ is an i-slice then $i = j = k$.

(Part B) Given 4 distinct points a, b, c, d such that $\{a, b\}$, $\{b, c\}$, $\{c, d\}$, and $\{a, d\}$ are i, j, k, and l-slices respectively then either $i = j = k = l$ or $i = k$ and $j = l$.

Step 2 If Ω is a k-slice then $f(\Omega)$ is a j-slice.

Step 3 If Ω and Ω' are k-slices then $f(\Omega)$ and $f(\Omega')$ are j-slices.

(Proof of Step 1 Part A) By definition of a k-slice if $i = j$ then $i = j = k$. Suppose $i \neq j$ then a and c differ in the i^{th} and j^{th} coordinate. But $\{a, c\}$ is a k-slice which is a contradiction.

(Proof of Step 1 Part B) Observe that since $\{a, b\}$ is a i-slice and $\{b, c\}$ is a j-slice the points a and c differ in the i^{th} and j^{th} coordinates. Similarly they differ in the l^{th} and k^{th} coordinate. Thus $\{i, j\} = \{k, l\}$. If $i = l$ then $i = j = k = l$ otherwise $i = k$ and $j = l$.

(Proof of Step 2) Any k-slice Ω is contained in a $k-$slice of the form

$$\Phi = \alpha_1 \times \ldots \alpha_{k-1} \times M_k \times \alpha_{k+1} \cdots \times \alpha_m$$

for $\alpha_i \in M_i$ so it suffices to show the result for Φ. Let $\eta \in M_k$ be an arbitrary point. Let δ_s and $\bar{\epsilon_\eta}$ be the uniquely geodesic constants for $\alpha_s \in M_s$ and $\eta \in M_k$ respectively. Define

$$W_\eta = \left(\prod_{s=1}^{k-1} B(\alpha_s, \epsilon_\eta)\right) \times B(\eta_t, \epsilon_\eta) \times \left(\prod_{s=k+1}^{m} B(\alpha_s, \epsilon_\eta)\right)$$

where $\epsilon_\eta = min\{\bar{\epsilon_\eta}, \delta_1, \dots, \delta_m\}$. Let $\Psi = W_\eta \cap \Phi$. For any three points $x_1, x_2, x_3 \in \Psi$ construct three isometrically embedded graphs $\iota_i : \mathcal{Q}^m_r \to W_\eta$ with $r = \frac{1}{4}d(x_i, x_{i+1})$, $\iota_i(e^r_1) = x_i$, and $\iota_i(-e^r_1) = x_{i+1}$ for $i \in \{1, 2, 3\}$ via Example 3.2 (note that $x_4 = x_1$). Since W_η is a product of uniquely geodesic metric spaces we can apply lemma 3.7 to $f \circ \iota_i$ to see that $\{f(x_1), f(x_2), f(x_3)\}$ satisfy the hypothesis of Step 1 part A. Thus $\{f(x_1), f(x_2), f(x_3)\}$ is a j-slice and since x_i were arbitrary $f(\Psi)$ is a j-slice. If Ψ' is a defined about a point $\eta' \in M_k$ sufficiently close to η then $f(\Psi) \cap f(\Psi')$ consists of more than two points and the set $f(\Psi) \cup f(\Psi')$ is a j-slice. This implies $f(\Phi)$ is a $j - slice$ since M_k has a single path component.

(Proof of Step 3) Note that

$$\Omega \subset \alpha_1 \times \cdots \times \alpha_{k-1} \times M_k \times \alpha_{k+1} \times \cdots \times \alpha_m = \Phi$$

and

$$\Omega' \subset \beta_1 \times \cdots \times \beta_{k-1} \times M_k \times \beta_{k+1} \times \cdots \times \beta_m = \Phi'$$

for some M_k and hence it suffices to show the result for Φ, Φ'. We can also assume that $\alpha_i = \beta_i$ for all $i \neq 1$ since we can change one coordinate at a time to interpolate between Φ and Φ'. Now there clearly exist distinct points $w_1, w_2 \in \Phi$ and $w'_1, w'_2 \in \Phi'$ such that $\{w_1, w'_1\}$ and $\{w_2, w'_2\}$ are 1-slices. By step 2 $f(w_1), f(w'_1), f(w_2), f(w'_2)$ satisfy the hypothesis of Step 1 part B. If $\{f(w_1), f(w'_1), f(w_2), f(w'_2)\}$ was a j-slice then by step 2 after applying the inverse isometry $\{w_1, w_2, w_1,' w'_2\}$ would be a k-slice or a 1-slice which is a contradiction. □

References

[1] M. Bridson and A. Haefliger. *Metric Spaces of Non-Positive Curvature.* Springer-Verlag, Berlin, 1999.

[2] Benson Farb and Howard Masur. *Teichmuller geometry of moduli space, I: Distance minimizing rays and the Deligne-Mumford compactification.* arXiv:0803.0141v1, 2008.

[3] W.W. Rouse Ball and H.S.M. Coxeter. *Mathematical Recreations and Essays.* Dover Publications, Inc., New York, thirteenth edition, 1987.

William Malone:
Dept. of Mathematics, University of Utah
155 S 1400 E Room 233,
Salt Lake City, UT 84112-0090
Email: malone@math.utah.edu